





Entomopathogenic fungi eria bassiana causes white muscardine disease ♦ Lubilosa project from 1990 – 2002: Metarhizium sp. Green muscle TM NOVACRID TM Missing field studies on adult individuals! ♦ Temperature preference of sporulation:

Night time: > 20°C, Day time: < 38°C

No oil controls conducted so far!



Stefan Jaronski - This image was released by the Agricultural Research Service, the research agency of the United States Department of Agriculture,

Neem oil (Azadirachta sp.)

- Schistocerca gregaria: decreased the flight performance substantially up to 70% (H. Schmutterer et al. Journal of Applied Entomology, 1993 https://doi.org/10.1111/j.1439-0418.1993.tb01186.x)
- Repellent effects of different products of the neem tree on the red locust, Nomadacris septemfasciata lasts 5 days (Langewald, et al. Pflanzenschutz, Umweltschutz 68, 55-57 1995. https://doi.org/10.1007/BF01996732)
- Strong phagorepellent effect on the desert locust, Schistocerca gregaria, and on the red locust, Nomadacris septemfasciata (Baumgart, et al. Journal of Applied Entomology. 116. 178 186. 2009. DOI 10.1111/j.1439-0418.1993.tb01186.x.)
- ♦ Neem oil product causes high mortality in flying S. gregaria (Nasseh, H. Wilps and S. KrallSource: Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz / Journal of PlantDiseases and Protection, Vol. 100, No. 6,1993. pp. 611-621

5

5

Hardening of linseed oil After mixing linseed with soda solution (b) 24 hours later

Screening for toxic essential oils

- Caraway, orange peel, wintergreen (birch oil), ginger, clove, peppermint, basil and Eucalypthus
- ♦ Highly toxic against *S. gregaria* and *L. migratoria*: caraway, orange peel and wintergreen oil in combination with the linseed oil emulsion.
- Publication: Abdelatti and Hartbauer 2019 Journal of Pest Science (//doi.org/10.1007/s1034 0-019-01169-7)
- ♦ Patent P18417pct: Formulation 102 and Formulation 103 (no orange peel oil)
- ♦ Has no toxic effects on adult mealworm beetles
- ♦ Moderate toxic for ladybug beetles
- ♦ Weat grass growth is mildly affected (some yellow tips after 3 weeks).

7

7







